

# Livestock Breeding and Genomics - Solution 3

Peter von Rohr

2018-10-12

## Problem 1: Breeding Value

We are considering a quantitative trait that depends on a given bi-allelic locus  $G$ . The frequency of the favorable allele corresponds to 0.08. Suppose that genotype frequencies follow the Hardy-Weinberg equilibrium. The difference between the homozygous genotypes corresponds to 15. The heterozygous genotype has a value of  $-1.5$ .

- Compute the breeding values and the dominance deviations for the three genotypes.
- Because of selecting the positive allele the frequency has increased to 0.096. How does this increased allele frequency change the breeding values?

**Hint:** Have a look at the summary table of all values in the course notes.

## Solution

According to the summary table of all values in the course notes, the breeding values depend on the term  $\alpha$ . Therefore, we start by computing  $\alpha$  first.

$$\alpha = a + (q - p)d$$

Based on the problem description, we know that  $a = 7.5$ ,  $p = 0.08$ ,  $q = 1 - p = 0.92$  and  $d = -1.5$ . Therefore

$$\alpha = 7.5 + (0.92 - 0.08) * (-1.5) = 6.24$$

- The summary table of all values then looks as follows.

Genotype	genotypic value	breeding value	dominance deviation
$G_iG_j$	$V_{ij}$	$BV_{ij}$	$D_{ij}$
$G_1G_1$	$a = 7.5$	$2q\alpha = 11.4816$	$-2q^2d = 2.5392$
$G_1G_2$	$d = -1.5$	$(q - p)\alpha = 5.2416$	$2pqd = -0.2208$
$G_2G_2$	$-a = -7.5$	$-2p\alpha = -0.9984$	$-2p^2d = 0.0192$

- Based on the change in the allele frequency to  $p = 0.096$  and  $q = 0.904$ . The value of  $\alpha$  changes to  $\alpha = 6.288$ . This has consequences for the whole summary table.

Genotype	genotypic value	breeding value	dominance deviation
$G_iG_j$	$V_{ij}$	$BV_{ij}$	$D_{ij}$
$G_1G_1$	$a = 7.5$	$2q\alpha = 11.368704$	$-2q^2d = 2.451648$
$G_1G_2$	$d = -1.5$	$(q - p)\alpha = 5.080704$	$2pqd = -0.260352$
$G_2G_2$	$-a = -7.5$	$-2p\alpha = -1.207296$	$-2p^2d = 0.027648$

Due to the increment in the allele frequency  $p$  from 0.08 to 0.096 the value of  $\alpha$  got bigger. But the breeding values decreased, because the negative influence of incrementing  $p$  on the breeding values was bigger than the positive change of  $\alpha$ .

## Problem 2: Allele Substitution

What is the meaning of the term **allele substitution** and how big is it in 1a) and 1b)?

### Solution

The effect of **allele substitution** occurs in the difference of the breeding values between two genotypes where one of these genotypes has one favorable allele more than the other. For a single bi-allelic locus there are two possible differences that fulfill the requirement from the previous sentence, namely  $BV_{12} - BV_{22}$  and  $BV_{11} - BV_{12}$ . The result of both differences is the same and corresponds to  $\alpha = a + (p - q) * d$ .

The allele substitution ( $\alpha$ ) in 1a) corresponds to 6.24 in 1b) the value is 6.288.

## Problem 3: Reading Data into R

You can download a file in csv-format from the course website. The URL is <https://charlotte-ngs.github.io/LBGFS2018/ex/w04/iris.csv>. Read the data from that csv-file into R using the function `read.csv2()`. Test the consequences of specifying the option `stringsAsFactors=FALSE`.

### Hints:

1. You can first download the csv-file to your local computer and then read the data, or you can directly indicate the URL when reading the data. You get more information with the command `?read.csv2` at the R-console.
2. Assign the result of `read.csv2()` to a variable
3. Use the function `str()` on the result of `read.csv2()` to see the difference between the two results of reading the data.
4. Use the description at <https://bookdown.org/rdpeng/rprogdatascience/getting-data-in-and-out-of-r.html> as a reference to read data into R. There is also a video on the same subject under [https://youtu.be/Z\\_dc\\_FADyi4](https://youtu.be/Z_dc_FADyi4).

### Solution

- Read the data directly using the URL:
- Read from the local computer

```
dfIris1 <- read.csv2(file = "iris.csv")
str(dfIris1)
```

```
## 'data.frame': 150 obs. of 5 variables:
## $ Sepal.Length: num 5.1 4.9 4.7 4.6 5 5.4 4.6 5 4.4 4.9 ...
## $ Sepal.Width : num 3.5 3 3.2 3.1 3.6 3.9 3.4 3.4 2.9 3.1 ...
## $ Petal.Length: num 1.4 1.4 1.3 1.5 1.4 1.7 1.4 1.5 1.4 1.5 ...
## $ Petal.Width : num 0.2 0.2 0.2 0.2 0.2 0.4 0.3 0.2 0.2 0.1 ...
## $ Species : Factor w/ 3 levels "setosa","versicolor",...: 1 1 1 1 1 1 1 1 1 1 ...
```

```
dfIris2 <- read.csv2(file = "iris.csv", stringsAsFactors = FALSE)
str(dfIris2)
```

```
## 'data.frame': 150 obs. of 5 variables:
## $ Sepal.Length: num 5.1 4.9 4.7 4.6 5 5.4 4.6 5 4.4 4.9 ...
## $ Sepal.Width : num 3.5 3 3.2 3.1 3.6 3.9 3.4 3.4 2.9 3.1 ...
## $ Petal.Length: num 1.4 1.4 1.3 1.5 1.4 1.7 1.4 1.5 1.4 1.5 ...
## $ Petal.Width : num 0.2 0.2 0.2 0.2 0.2 0.4 0.3 0.2 0.2 0.1 ...
## $ Species : chr "setosa" "setosa" "setosa" "setosa" ...
```

### Additional Problem: Create a plot in R

Plot the values in the columns `Sepal.Length` and `Petal.Length` of the Iris data set. The plot should look like the following figure.

### Solution

```
plot(dfIris2$Sepal.Length, dfIris2$Petal.Length)
```

