

# Livestock Breeding and Genomics - Exercise 3

*Peter von Rohr*

*2019-10-11*

## Problem 1: Breeding Value

We are considering a quantitative trait that depends on a given bi-allelic locus  $G$ . The frequency of the favorable allele corresponds to 0.08. Suppose that genotype frequencies follow the Hardy-Weinberg equilibrium. The difference between the homozygous genotypes corresponds to 15. The heterozygous genotype has a value of  $-1.5$ .

- a) Compute the breeding values and the dominance deviations for the three genotypes.
- b) Because of selecting the positive allele the frequency has increased to 0.096. How does this increased allele frequency change the breeding values?

**Hint:** Have a look at the summary table of all values in the course notes.

## Problem 2: Allele Substitution

What is the meaning of the term **allele substitution** and how big is it in 1a) and 1b)?

## Problem 3: Reading Data into R

You can download a file in csv-format from the course website. The URL is <https://charlotte-ngs.github.io/LBGFS2019/ex/w04/iris.csv>. Read the data from that csv-file into R using the function `read.csv2()`. Test the consequences of specifying the option `stringsAsFactors=FALSE`.

**Hints:**

1. You can first download the csv-file to your local computer and then read the data, or you can directly indicate the URL when reading the data. You get more information with the command `?read.csv2` at the R-console.
2. Assign the result of `read.csv2()` to a variable
3. Use the function `str()` on the result of `read.csv2()` to see the difference between the two results of reading the data.
4. Use the description at <https://bookdown.org/rdpeng/rprogdatascience/getting-data-in-and-out-of-r.html> as a reference to read data into R. There is also a video on the same subject under [https://youtu.be/Z\\_dc\\_FADyi4](https://youtu.be/Z_dc_FADyi4).

## Additional Problem: Create a plot in R

Plot the values in the columns `Sepal.Length` and `Petal.Length` of the Iris data set.