

# Livestock Breeding and Genomics - Solution 6

Peter von Rohr

2022-10-28

## Problem 1: Regression Model

Use the dataset on weaning weight and fit a regression model of weaning weight on breast circumference. The following tasks are to be completed.

- Use matrix-vector notation to specify the model and fill the information from the dataset into the model.
- Compute the solution for the estimated regression coefficient using a least squares approach.
- Use R to verify your result.

The data set is available from

```
## https://charlotte-ngs.github.io/lbgfs2022/data/beef\_data\_bc.csv
```

## Solution

The regression model in matrix-vector notation is given by the following formula

$$y = X\beta + e$$

where  $y$  is the vector of known responses. For our example this is the vector of weaning weights. The vector  $\beta$  contains the intercept and the regression coefficient. The vector  $e$  contains the random residuals. The matrix  $X$  contains a column of all ones and a second column with breast circumference.

The data are imported to a tibble using

```
tbl_ww <- readr::read_delim(file = s_data_path, delim = ",",
                           col_types = readr::cols(
                             Animal = readr::col_integer(),
                             Sire = readr::col_integer(),
                             Dam = readr::col_integer(),
                             Herd = readr::col_factor(),
                             `Weaning Weight` = readr::col_double(),
                             `Breast Circumference` = readr::col_double()
                           ))
head(tbl_ww)
```

```
## # A tibble: 6 x 6
##   Animal Sire  Dam Herd `Weaning Weight` `Breast Circumference`
##   <int> <int> <int> <fct>          <dbl>                <dbl>
```

```
## 1      12      1      4 1          2.61          1.62
## 2      13      1      4 1          2.31          1.96
## 3      14      1      5 1          2.44          1.48
## 4      15      1      5 1          2.41          1.47
## 5      16      1      6 2          2.51          1.5
## 6      17      1      6 2          2.55          1.47
```

The information for the known vector  $y$  and for matrix  $X$  are taken from the dataset. The vector  $y$  contains all weaning weight values

```
vec_y <- tbl_ww$`Weaning Weight`
```

The matrix  $X$  is a matrix with two columns and as many rows as there are observations.

```
n_nr_obs <- nrow(tbl_ww)
mat_X <- matrix(c(rep(1,n_nr_obs), tbl_ww$`Breast Circumference`), nrow = n_nr_obs, ncol = 2)
```

The least squares estimate is

$$\hat{\beta} = (X^T X)^{-1} X^T y$$

Inserting the numbers into the formula leads to

```
mat_xtx <- crossprod(mat_X)
mat_xty <- crossprod(mat_X, vec_y)
vec_hat_beta <- solve(mat_xtx, mat_xty)
vec_hat_beta
```

```
##           [,1]
## [1,] 1.638243
## [2,] 0.561967
```

The standard deviation of the residuals is used as an estimate of the spread of the error terms.

```
vec_r <- vec_y - crossprod(t(mat_X), vec_hat_beta)
n_sd_res <- sqrt(crossprod(vec_r) / (n_nr_obs-2))
n_sd_res
```

```
##           [,1]
## [1,] 0.315731
```

Verification of the results using R

```
lm_ww <- lm(`Weaning Weight` ~ `Breast Circumference`, data = tbl_ww)
summary(lm_ww)
```

```
##
## Call:
## lm(formula = `Weaning Weight` ~ `Breast Circumference`, data = tbl_ww)
##
```

```
## Residuals:
##      Min       1Q   Median       3Q      Max
## -0.45921 -0.15326  0.01814  0.10909  0.61699
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)         1.6382     0.7775   2.107  0.0536 .
## 'Breast Circumference' 0.5620     0.5019   1.120  0.2817
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.3157 on 14 degrees of freedom
## Multiple R-squared:  0.0822, Adjusted R-squared:  0.01664
## F-statistic: 1.254 on 1 and 14 DF,  p-value: 0.2817
```

## Problem 2: Fixed Linear Effects Model

Use the same dataset as in Problem 1 and fit a fixed linear effects model using breast circumference and herd as fixed effects in a model. Use the same path to get to the solution as in Problem 1 and complete the same set of tasks.

### Solution

The model in matrix vector notation is the same as in Problem 1

$$y = X\beta + e$$

But the definition of the vector  $\beta$  and the matrix  $X$  are different. The vectors  $y$  and  $e$  are defined the same way.

The check with R

```
lm_ww_bc_herd <- lm(`Weaning Weight` ~ `Breast Circumference` + Herd, data = tbl_ww)
summary(lm_ww_bc_herd)
```

```
##
## Call:
## lm(formula = `Weaning Weight` ~ `Breast Circumference` + Herd,
##     data = tbl_ww)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -0.50908 -0.12643  0.00342  0.10719  0.57491
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)         1.48141     0.88203   1.680  0.117
## 'Breast Circumference' 0.63990     0.54870   1.166  0.264
## Herd2                 0.07344     0.17260   0.426  0.677
##
## Residual standard error: 0.3254 on 13 degrees of freedom
## Multiple R-squared:  0.0948, Adjusted R-squared: -0.04446
## F-statistic: 0.6808 on 2 and 13 DF,  p-value: 0.5234
```