

Livestock Breeding and Genomics - Exercise 2

Peter von Rohr

2022-09-29

Problem 1: Vectors in R

Vector Definition

Although there exists a function called `vector()` in R, vectors are always defined in R using the function `c()` which stands for “concatenation”.

Vector Assignment

Let us assume we want to assign the following vector a

$$a = \begin{bmatrix} 10 \\ 7 \\ 43 \end{bmatrix}$$

to the variable named `a` in R, then this can be done with the following statement

```
a <- c(10,7,43)
```

Access of single Vector Element

A single vector element can be accessed using the variable name followed by the element index in brackets. Hence, if we want to know the first element of vector `a`, we have to write

```
a[1]
```

```
## [1] 10
```

Computations with Vector Elements

Vector elements can be used in arithmetic operations such as summation, subtraction and multiplication as shown below

```
a[1] + a[3]
```

```
## [1] 53
```

```
a[2] * a[3]
```

```
## [1] 301
```

```
a[3] - a[1]
```

```
## [1] 33
```

The function `sum()` can be used to compute the sum of all vector elements. The function `mean()` computes the mean of all vector elements.

```
sum(a)
```

```
## [1] 60
```

```
mean(a)
```

```
## [1] 20
```

Vector Computations

Arithmetic operations can also be performed not only on elements of vectors but also on complete vectors. Hence, we can add the vector `a` to itself or we can multiply it by a factor of 3.5 which is shown in the following code-chunk

```
a + a
```

```
## [1] 20 14 86
```

```
3.5 * a
```

```
## [1] 35.0 24.5 150.5
```

More Computations on Vectors

Given are the following two vectors v and w .

$$v = \begin{bmatrix} 3 \\ -5 \\ 1 \\ 9 \end{bmatrix}$$

$$w = \begin{bmatrix} 1 \\ 9 \\ -12 \\ 27 \end{bmatrix}$$

Compute

- the sum $v + w$,
- the difference $v - w$ and
- the dot product $v \cdot w$.

Problem 2: Matrices in R

Matrices in R are defined using the function `matrix()`. The function `matrix()` takes as first arguments all the elements of the matrix as a vector and as further arguments the number of rows and the number of columns. The following statement generates a matrix with 4 rows and 3 columns containing all integer numbers from 1 to 12.

```
mat_by_col <- matrix(1:12, nrow = 4, ncol = 3)
mat_by_col
```

```
##      [,1] [,2] [,3]
## [1,]   1   5   9
## [2,]   2   6  10
## [3,]   3   7  11
## [4,]   4   8  12
```

As can be seen, the matrix elements are ordered by columns. Often, we want to define a matrix where elements are filled by rows. This can be done using the option `byrow=TRUE`

```
mat_by_row <- matrix(1:12, nrow = 4, ncol = 3, byrow = TRUE)
mat_by_row
```

```
##      [,1] [,2] [,3]
## [1,]   1   2   3
## [2,]   4   5   6
## [3,]   7   8   9
## [4,]  10  11  12
```

Access of Matrix Elements

Matrix elements can be accessed similarly to what was shown for vectors. But to access a single element, we need two indices, one for rows and one for columns. Hence the matrix element in the second row and third column can be accessed by

```
mat_by_row[2,3]
```

```
## [1] 6
```

Arithmetic Computations with Matrices

Arithmetic computations with matrices can be done with the well-known operators as long as the matrices are compatible. For summation and subtraction matrices must have the same number of rows and columns. For matrix-multiplication, the number of columns of the first matrix must be equal to the number of rows of the second matrix.

In R the arithmetic operators `+`, `-` and `*` all perform element-wise operations. The matrix multiplication can either be done using the operator `%*%` or the function `crossprod()`. It has to be noted that the statement

```
crossprod(A, B)
```

computes the matrix-product $A^T \cdot B$ where A^T stands for the transpose of matrix A . Hence the matrix product $A \cdot B$ would have to be computed as

```
crossprod(t(A), B)
```

More Examples

Given the matrices X and Y

```
X <- matrix(1:15, nrow = 5, ncol = 3)
Y <- matrix(16:30, nrow = 5, ncol = 3)
```

Compute

- $X + Y$
- $Y - X$
- multiplication of elements between X and Y
- matrix-product $X^T \cdot Y$
- matrix-product $X^T \cdot X$
- matrix-product $Y^T \cdot Y$